FACS Buffer

Dillon Corvino

2026-09-18

Protocol ID: BUF-FACS-BUF **Version:** v1.0 **Author:** Dillon Corvino **Date:** 2026-09-18

Purpose

This buffer is used for washing, resuspending, and handling cells during **flow cytometry staining and sorting (FACS)**. It contains no EDTA, unlike MACS Buffer (`BUF-MACS-BUF`), so it is the correct choice for perfusion and for any step upstream of an EDTA-sensitive assay (e.g. collagenase digestion).

Table of contents

Purpose 1

Linked protocols 1

Working buffer composition 2

Preparation 2

A. Preparation of stock (if applicable) 2

B. Preparation of working buffer 2

Storage and stability 2

Reagent details 2

Safety (brief) 3

Version history 3

Linked protocols

- `MUS-SPL-DIS`
- `MUS-LIV-DIG`
- `MUS-LNG-DIG`
- `MUS-BM-ISO`

- `MUS-BLD-PRC`
- `MUS-TMR-DIG`

Working buffer composition

Typical preparation for **500 mL**:

Component	Stock concentration	Volume for 500 mL	Final concentration	Notes
PBS 1x	—	490 mL	—	Base buffer
FCS	—	10 mL	2% (v/v)	Heat-inactivated if required

> **Note**
>
> FACS Buffer and MACS Buffer (`BUF-MACS-BUF`) share the same PBS/FCS base — MACS Buffer additionally contains 2 mM EDTA. See `BUF-MACS-BUF` for that recipe.

Preparation

A. Preparation of stock (if applicable)

Not applicable — prepared directly as the working buffer.

B. Preparation of working buffer

1. Start with sterile **PBS 1x** in an appropriate bottle.
2. Add **FCS** to achieve **2% (v/v)**.
3. Mix gently by inversion.
4. If sterility is required, filter through a **0.22 um filter** into a sterile storage bottle.
5. Label clearly with buffer name, date of preparation, initials, and sterile/non-sterile status.

Storage and stability

- Store at **4 °C**.
- Recommended use within **4–6 weeks**, subject to internal QC and sterility checks.
- Invert gently before use to resuspend any settled components.
- Avoid repeated temperature cycling; prepare smaller working aliquots if accessed frequently.

Reagent details

Component	Supplier	Cat#	Notes
PBS 1x	[TBD]	[TBD]	Sterile, Ca2+/Mg2+-free recommended
FCS	[TBD]	[TBD]	Heat-inactivated if required

Safety (brief)

- Handle biological samples in FACS buffer according to institutional biosafety rules.
- FCS is of animal origin and should be treated as a potential biohazard.

Version history

Version	Date	Author	Change summary
v1.0	2026-09-18	Dillon Corvino	Split out of `BUF-FACS-MACS` (formerly `buf_facs_macs_buffer.qmd`) so each buffer_id maps to exactly one recipe.